



COMMUNICATION

ON THE MAXIMUM NUMBER OF EDGES IN A HYPERGRAPH WHOSE LINEGRAPH CONTAINS NO CYCLE

J.-C. BERMOND

Université Paris-Sud, Laboratoire de Recherche en Informatique, Bât. 490, 91405 Orsay, France

P. FRANKL

C.M.S., 54 Bd Raspail, 75006 Paris, France

F. STERBOUL

Université de Lille, France

Communicated by C. Berge
Received 14 January 1980

Soit $H = (X, \mathcal{E})$ un hypergraphe h -uniforme avec $|X| = n$ et soit $L_{h-1}(H)$ le graphe dont les sommets représentent les arêtes de H , deux sommets étant reliés si et seulement si les arêtes qu'ils représentent intersectent en $h-1$ sommets. Nous montrons que si $L_{h-1}(H)$ ne contient pas de cycle, alors $|\mathcal{E}| < \binom{n}{h-1}/h-1$, la borne étant exacte pour $h=2$ et pour des valeurs de n pour $h=3$. Ce problème mène à une conjecture sur les "presque systèmes de Steiner".

Let $H = (X, \mathcal{E})$ be a h -uniform hypergraph, with $|X| = n$ and let $L_{h-1}(H)$ be the graph, whose vertices are the edges of H , two vertices being joined if and only if the edges they represent intersect in $h-1$ vertices. We prove that, if $L_{h-1}(H)$ contains no cycle, then $|\mathcal{E}| < \binom{n}{h-1}/h-1$; moreover the bound is exact for $h=2$ and with some values of n for $h=3$. This problem leads to a conjecture on "almost Steiner systems".

1. Introduction

Let X be a set of n elements and let $H = (X, \mathcal{E})$ be a h -uniform hypergraph, that is \mathcal{E} is a family of h -subsets of X .

We denote by $L_{h-1}(H)$ the graph, whose vertices are the edges of H and where $E_1, E_2 \in \mathcal{E}$ are joined by an edge iff $|E_1 \cap E_2| = h-1$. These graphs were previously studied (see for references [2]).

In this note we answer the following problem, which is a version of a problem due to Kachelsky and Kleitman, presented at the Problem Session of the 7th British Combinatorial Conference (Cambridge, August 1979).

Theorem 1. Suppose $L_{h-1}(H)$ contains no cycle, then

$$|\mathcal{E}| \leq \frac{\binom{n}{h-1}}{h-1}. \quad (1)$$

Moreover the bound is exact for $h=2$ and for some values of n for $h=3$.

2. Proof of inequality (1)

Let $F \subset X$, $|F| = h-1$. The F is contained in at most 2 different members of \mathcal{E} , as otherwise for $E_1, E_2, E_3 \in \mathcal{E}$ and containing F , we would have a triangle in $L_{h-1}(H)$.

Hence, for $F \subset E \in \mathcal{E}$, $|F| = h-1$ there are only two possibilities, either F is contained only in E or there is exactly one more edge E' , with $F \subset E'$.

Let $a(E)$ denote the number of $(h-1)$ -subsets of $E \in \mathcal{E}$, which are contained in some $E' \in \mathcal{E}$, $E' \neq E$. Then the vertex E is joined to exactly $a(E)$ other vertices of $L_{h-1}(H)$.

Let m be the number of edges of $L_{h-1}(H)$, then we obtain

$$2m = \sum_{E \in \mathcal{E}} a(E). \quad (2)$$

On the other hand, $L_{h-1}(H)$ contains no cycle, hence we also have

$$m \leq |\mathcal{E}| - 1. \quad (3)$$

Combining (2) and (3) we obtain

$$|\mathcal{E}| - 1 \geq \frac{1}{2} \sum_{E \in \mathcal{E}} a(E). \quad (4)$$

Let $c(H)$ be the number of $(h-1)$ -subsets of X , which are contained in some edge E of H . Obviously $c(H) \leq \binom{n}{h-1}$. We obtain

$$c(H) = \sum_{E \in \mathcal{E}} (h - a(E)) + \frac{1}{2} \sum_{E \in \mathcal{E}} a(E) \leq \binom{n}{h-1}$$

or equivalently

$$c(H) = h|\mathcal{E}| - \frac{1}{2} \sum_{E \in \mathcal{E}} a(E) \leq \binom{n}{h-1}. \quad (5)$$

From (5) using (4) we deduce

$$h|\mathcal{E}| - (|\mathcal{E}| - 1) = (h-1)|\mathcal{E}| + 1 \leq \binom{n}{h-1}$$

or equivalently $|\mathcal{E}| \leq \binom{n}{h-1}/(h-1)$ proving (1).

Remark. Let us define $L_{\leq i}(H)$ to be the graph whose vertices are the edges of H and where $E_1 \neq E_2 \in \mathcal{E}$ are joined iff $|E_1 \cap E_2| \geq i$ ($1 \leq i \leq h-1$). Suppose that

$L_{\infty i}(H)$ contains no cycle. Then the above proof yields

$$|\mathcal{E}| < \binom{n}{i} / \left(\binom{h}{i} - 1 \right).$$

3. Constructions

For $h = 2$, we can take any tree to show that the bound (1) is exact.

Let now $h = 3$.

Suppose $n = 8k + 5$ is a prime and α is a primitive root modulo n , that is the numbers $\pm 2^\alpha$, $\alpha = 0, 1, \dots, \frac{1}{2}(n-3)$ are all different. (It is conjectured in number theory that there exists an infinite number of such n 's.)

Let H be a hypergraph with vertex set $\{0, 1, \dots, n-1\}$ and edge set:

$$\begin{aligned} \{E_i^\alpha = \{i, i + 2^{2\alpha}, i + 2^{2\alpha+1}\}, i = 1, \dots, n-1; \alpha = 0, 1, \dots, \tfrac{1}{4}(n-5)\} \\ \cup \{E_0^\alpha = \{0, 2^{2\alpha+1}, 2^{2\alpha+2}\}, \alpha = 0, 1, \dots, \tfrac{1}{4}(n-9)\} \end{aligned}$$

(all the numbers are taken modulo n).

We assert that $L_{h-1}(H)$ contains no cycle. To see this, observe that the subgraph generated by $X^\alpha = \{E_i^\alpha, i = 1, \dots, n-1\}$ is a Hamiltonian path on X^α .

Moreover $|E_i^\alpha \cap E_j^{\alpha'}| \leq 1$, for $1 \leq i, j \leq n-1$, $\alpha \neq \alpha'$ as 2 is a primitive root. Therefore there are no edges in $L_{h-1}(H)$ joining X^α to $X^{\alpha'}$.

Finally the vertex E_0^α is joined in $L_{h-1}(H)$ only to $E_{2^{2\alpha+1}}^\alpha$ and to $E_{2^{2\alpha+2}}^\alpha$. Hence $L_{h-1}(H)$ is a tree (see Fig. 1 for an example, $n = 13$).

We have $|\mathcal{E}| = \frac{1}{4}(n-1)(n-1) + \frac{1}{4}(n-5) = \frac{1}{4}n(n-1) - 1$ which proves that the bound (1) is optimal for the mentioned values of n .

For the other values of n , we can prove:

Theorem 2. *For every n , there exists a 3-uniform hypergraph H of order n such that $L_2(H)$ has no cycle and $\frac{1}{2}\binom{n}{3} - |H| \leq \frac{1}{4}n$.*

We use the following lemma:

Lemma 1. *There exists a sequence of matchings (graphs with maximum degree 1) C_j such that*

- (a) $|C_{2i-1}| = |C_{2i}| = i$,
- (b) *the vertex-set of $G_j = C_1 \cup \dots \cup C_j$ is $\{x_1, \dots, x_{j+1}\}$*
- (c) $C_j \cap C_k = \emptyset$ ($j \neq k$),
- (d) *the degree-sequence of G_{2i} is dominated by $\{1, 2, 3, \dots, 2i-3, 2i-2, 2i-1, 2i-1\}$, for $i \geq 3$.*

The construction of H in Theorem 2 is the following:

$$H = \{\{x_j\} \cup E \mid E \in C_{j-2}, j = 3, \dots, n\}.$$




Fig. 1.

Lemma 1 is proved by induction, with the aid of the Tutte-Berge matching theorem (see for instance [1]). This method is also valid for $h > 3$. However, in this case, Lemma 1 should be replaced by results not yet proved such as the following one:

Conjecture. There exists a sequence of $(h-1)$ -uniform hypergraphs S_i such that:

- (a) the vertex-set of $S_1 \cup \dots \cup S_i$ is $\{x_1, \dots, x_{i+h-2}\}$,
- (b) $S_i \cap S_k = \emptyset$ ($j \neq k$),
- (c) every two edges of S_i meet in at most $h-3$ vertices,
- (d) $|S_i| = \binom{i+h-2}{h-2} / (h-1) - O(j^{h-3})$.

The last two conditions mean that S_i is an "almost Steiner system". (Note $O(j^{h-3}) \leq c_h j^{h-3}$ for some constant c_h).

References

- [1] C. Berge, Graphs and hypergraphs (North-Holland, Amsterdam, 1976).
- [2] J.-C. Bermond, A. Germa and M.-C. Heydemann, Graphes représentatifs d'hypergraphes, in: Coll. Math. Discretes, Bruxelles (1978), Cahiers du C.E.R.O. 20 (1978) 325-329.